

Pediatric Behavioral Health Medication Initiative Prior Authorization (PA) Request Form

Please fax form to **617.673.0988** or mail to Tufts Health Plan, 1 Wellness Way, Canton, MA 02021-1166, Attn: Pharmacy Utilization Management Department. For online prior authorization: https://point32health.promptpa.com. *Complete this form for Tufts Health Together (MassHealth) only.*

Member In	formation				
Member Last Name:		Memb	Member First Name:		
Member ID: DOB:		DOB:	l	Gender: [□Male □Female □Unknown
				1	
Prescriber	Information				
Prescriber	Last Name:		Prescrib	er First Nan	ne:
Specialty: Phone #:		Phone #:			Secure Fax #:
NPI #:			DEA/xDI	ĒA:	
Prescriber	Point of Contact Name	(POC) (if different t	han provider	·):	
POC Email	(not required):				
Prescriber					
	Representative Signat	ure:			
Date:					
/ledical Inf		an acuto caro cottin	, a 2		
	e member currently in : □ Yes (Inpatient)	an acute care settii	ıgı		
	☐ Yes (Community Ba	sad Acuta Traatma	nt)		
	☐ Yes (Partial Hospita		1110)		
		1124110117			
		acute care setting	nlease docu	iment outp	atient prescriber after discharge.
	☐ Prescriber name:	-	•	·	•
	☐ Contact information				
	the member been hosp		iatric conditi	on within t	he past three months?
	☐ Yes. Please docume				
	 □ No				
4. Is the member considered a severed risk of harm to self or others?					

		Yes. Please provide details:
		No.
5.	Dlease	No adjects the prescriber's specialty
э.		ndicate the prescriber's specialty
	_	Psychiatry
		Neurology Other
	ш	Specialist consult details (if prescriber submitting request is not a specialist)
		□ Name(s) of specialist(s):
		Specialty:
		Date(s) of last visit or consult:
_	-	☐ Contact information:
6.		l-level practitioners (e.g., nurse practitioners, physician assistants), please provide the name and specialty
_		borating physician:
7.		document member custody status:
		Parent/Guardian
_	⊔	Department of Children and Families (DCF)
8.		document member placement status:
		Home with Parent/Guardian
		Foster Care
		Residential Treatment Facility
		Uncertain
		Other:
9.	Please	document agency involvement:
		Department of Children and Families (DCF)
		Department of Mental Health (DMH)
		Department of Developmental Services (DDS)
		Department of Youth Services (DYS)
10.		nember currently receiving appropriate psychotherapeutic and/or community-based services for the
	_	d clinical mental health concerns (e.g., Applied Behavioral Analysis, Children's Behavioral Health
	Initiativ	e, school interventions, specialized placement)?
		Yes. Please document details of intervention, if applicable:
		No No
11		tric care provided is coordinated with other psychotherapeutic and community-based services.
11.	•	Yes
		No
12		nember a referral candidate for care coordination?
12.		Yes
		No No
12		
13.	benefic	nswer to number 12 is "yes", please describe which additional behavioral health services would be ial.

14. Please provide the member's complete treatment regimen (including all current and requested behavioral health medications). Include each medication's name, dose, frequency, and indication.

Medication name/dose/frequency	Indication
1.	1.
2.	2.
3.	3.
4.	4.
5.	5.
6.	6.
7.	7.

POLYPHARMACY

Antidepressant Polypharmacy

Complete this section for all members less than 18 years of age if the request will result in antidepressant polypharmacy (ove

erla _l	pping pl	narmacy claims for two or more antidepressants for at least 60 days within a 90 day period).
15.	For ant	idepressant polypharmacy, please indicate whether one of the following applies: Member is cross titrating/tapering antidepressant therapy Member had an inadequate response (defined as four weeks of therapy) or adverse reaction to two monotherapy trials as clinically appropriate Antidepressant polypharmacy regimen of two or fewer antidepressants includes one of the following agents: bupropion, mirtazapine, or trazodone One antidepressant in the regimen is indicated for a comorbid condition in which antidepressants may be clinically appropriate
16.		document if monotherapy trials (include drug name, dates/duration of use, and outcome) were tried prescribing polypharmacy with two or more antidepressants for this member.
17.	Please	document clinical rationale for antidepressant polypharmacy for this member.
18.		document the treatment plans for medication regimen simplification (e.g., dose consolidation, frequency on) or medical necessity for continuation of a complex medication regimen.

Benzodiazepine Polypharmacy

Complete this section for all members less than 18 years of age if the request will result in benzodiazepine polypharmacy (overlapping pharmacy claims for two or more benzodiazepines for at least 60 days within a 90 day period).

19.	Does the member have a seizure diagnosis only?
	□ Yes
	□ No
20.	Is the member cross-titrating/tapering benzodiazepine therapy?
	□ Yes
	□ No
21.	Please document if monotherapy trials (include drug name, dates/duration of use, and outcome) were tried
	before prescribing polypharmacy with two or more benzodiazepines for this member.
22.	Please document clinical rationale for benzodiazepine polypharmacy for this member.
23.	Please document the treatment plans for medication regimen simplification (e.g., dose consolidation, frequency
	reduction) or medical necessity for continuation of a complex medication regimen.
Cerebr	ral Stimulant Polypharmacy
Comple	ete this section for all members less than 18 years of age if the request will result in cerebral stimulant
polypha	armacy (overlapping pharmacy claims for two or more cerebral stimulants for at least 60 days within a 90 day
period;	immediate-release and extended-release formulations of the same chemical entity are counted as one).
24.	For cerebral stimulant polypharmacy, please select all that apply:
	☐ Member had an inadequate response (defined as more than 7 days of therapy), adverse reaction or
	contraindication to monotherapy trial with a methylphenidate product Member had an inadequate response (defined as more than 7 days of therapy), adverse reaction or
	contraindication to monotherapy trial with an amphetamine product
25	Please document any additional monotherapy trials (include drug name, dates/duration of use, and outcome)
23.	that were tried before prescribing polypharmacy with two or more cerebral stimulants for this member.
26	Please document clinical rationale for cerebral stimulant polypharmacy for this member.
20.	

27.		document the treatment plans for medication regimen simplification (e.g., dose consolidation, frequency on) or medical necessity for continuation of a complex medication regimen.				
Mood	Stabilize	er Polypharmacy				
-		ection for all members less than 18 years of age if the request will result in mood stabilizer polypharmacy armacy claims for three or more mood stabilizers for at least 60 days within a 90 day period).				
28.	Does th	e member have a seizure diagnosis only?				
		Yes				
29.		elect all that apply:				
		Member has a diagnosis for which mood stabilizers may be clinically appropriate (e.g., migraine, neuropathic pain)				
		Member has tried and failed other clinically appropriate therapies				
		Member is cross-titrating or tapering mood stabilizers therapy				
		Member has had an inadequate response or adverse reaction to two monotherapy trials and/or combination therapy trials as clinically appropriate				
30.	Please document if monotherapy trials (include drug name, dates/duration of use, and outcome) were tried before prescribing polypharmacy with three or more mood stabilizers.					
31.	Please o	document clinical rationale for mood stabilizer polypharmacy.				
32.		document the treatment plans for medication regimen simplification (e.g., dose consolidation, frequency on) or medical necessity for continuation of a complex medication regimen.				
Antips	ychotic	Polypharmacy				
•		ection for all members less than 18 years of age if the request will result in antipsychotic polypharmacy armacy claims for two or more antipsychotics for at least 60 days within a 90 day period).				
33.	Does th place?	e member have a comprehensive behavioral health plan (i.e., non-pharmacologic interventions) in				
		Yes				
		No				
34.	Please s	elect the stage of treatment and clinical rationale for antipsychotic polypharmacy.				
		Acute Stage (defined as initiation of antipsychotic treatment likely with subsequent dose adjustments to maximize response and minimize side effects)				

	Medication Name/Dose/Frequency	Dates/Duration of Use			
	1.	1.			
	2.	2.			
		Member is transitioning from one antipsychotic to the other. Other, please explain:			
□ Main	tenance Stage (response to antipsychotic treatm	ent with goal of remission or recovery)			
	s the regimen effective, therapy benefits outweig]Yes]No	h risks, and appropriate monitoring is in place?			
	resulted in symptom exacerbation	_			
	Other significant barrier for antipsychotic	therapy discontinuation. Please explain:			
□ Disco taper	Other significant barrier for antipsychotic No Indicated that the red); select all that apply:	antipsychotic regimen can likely be successfully			
□ Disco taper	Other significant barrier for antipsychotic No Continuation Stage (clinically indicated that the red); select all that apply: Member is transitioning from one antipsychotic	antipsychotic regimen can likely be successfully			
□ Disco taper	Other significant barrier for antipsychotic No Portinuation Stage (clinically indicated that the red); select all that apply: Member is transitioning from one antipsychotic. Member is tapering antipsychotic. Please desc	antipsychotic regimen can likely be successfully			

Multiple

Complete behaviora

35. Please document monotherapy trials (include drug name, dates/duration of use, and outcome) tried before prescribing a polypharmacy regimen for this member.

Medication Name	Dates/Duration of Use	Outcomes
1.	1.	1.
2.	2.	2.

	3.	3.	3.
	4.	4.	4.
	5.	5.	5.
	6.	6.	6.
	7.	7.	7.
36	Does the member have a seizur	e diagnosis only?	
50.	☐ Yes	and the state of t	
	□ No		
37.	Please select all that apply:		
	, , ,	rating or tapering mood stabilizer	
			action to two monotherapy trials and/or
		on trials as clinically appropriate	.,
	☐ Member has a diag	nosis in which mood stabilizers may b	oe clinically appropriate (e.g., migraine, ly appropriate therapies have been tried
		niatric and comorbid diagnosis in wh	ich mood stabilizers may be clinically
		igraine, neuropathic pain) and docur	mentation that other clinically relevant
38.	Please document the clinical rat	ionale for use of multiple behavioral	health medications for this member who is
_	less than 18 years of age.		
-			
39.		plans for medication regimen simpli for continuation of a complex medic	ification (e.g., dose consolidation, frequency ation regimen.

AGE LIMIT

Medication Name

Request for One of the Following Classes of Medications for Members Less than 6 Years of Age: Antidepressant, Atomoxetine, Benzodiazepine, Buspirone, or Mood Stabilizer

Please document any previous medication trial(s). Include drug name, dates/duration of use, and outcome.

Dates/Duration of Use

Outcomes

_		_	_
1.		1.	1.
2.		2.	2.
2.		2.	۷.
3.		3.	3.
4.		4.	4.
4.		4.	4.
5.		5.	5.
6.		6.	6.
7.		7.	7.
	stabilizer for this member w		
st for an	n Antipsychotic for Member	rs Less Than 6 Years of Age	
. Does t	he member have a comprel	nensive behavioral treatment plan (i.e.,	non-pharmacological intervention
place?	·	• , ,	
. \square	Yes		
	NI -		
	No		
	select the stage of treatme	nt and clinical rationale for antipsychot	, ,,
	select the stage of treatme Acute Stage (initiation of	antipsychotic treatment likely with sub	, ,,
	select the stage of treatme Acute Stage (initiation of response and minimize side)	antipsychotic treatment likely with sub de effects)	sequent dose adjustments to maxi
	select the stage of treatme Acute Stage (initiation of response and minimize sid Maintenance Stage (respo	antipsychotic treatment likely with sub de effects) onse to antipsychotic treatment with go	sequent dose adjustments to maxioal of remission or recovery)
	select the stage of treatme Acute Stage (initiation of response and minimize sid Maintenance Stage (response in the regimen effective)	antipsychotic treatment likely with sub de effects) onse to antipsychotic treatment with go ve, therapy benefits outweigh risks, and	sequent dose adjustments to maxioal of remission or recovery)
	select the stage of treatme Acute Stage (initiation of response and minimize sid Maintenance Stage (response in the regimen effective in Yes	antipsychotic treatment likely with sub de effects) onse to antipsychotic treatment with go oe, therapy benefits outweigh risks, and	sequent dose adjustments to maxional of remission or recovery)
	select the stage of treatme Acute Stage (initiation of response and minimize sid Maintenance Stage (response in the regimen effective in the stage) Has the member been	antipsychotic treatment likely with sub de effects) onse to antipsychotic treatment with go ve, therapy benefits outweigh risks, and	sequent dose adjustments to maxical of remission or recovery) d appropriate monitoring is in place ths or longer?

	Family/caregi	mptom exacerbation ver does not support the antipsychotic rbation ant barrier for antipsychotic therapy dis	
	tapered) Member is transiti	(clinically indicated that the antipsychooning from one antipsychotic to another gantipsychotic. Please describe taper p	er
•	•	Cerebral Stimulant for Members Less tedication trial(s). Include drug name, da	· · · · · · · · · · · · · · · · · · ·
1.	Medication Name	Dates/Duration of Use 1.	Outcomes 1.
1.		1.	1.
2.		2.	2.
3.		3.	3.
4.		4.	4.
5.		5.	5.
6.		6.	6.
7.		7.	7.
Please d	e member have a cardiova Yes No document clinical rationale n three years of age.	ascular diagnosis only? e for use of an alpha ₂ agonist and/or cer	rebral stimulant for this member wh

Request for a Hypnotic Medication for Members Less than 6 Years of Age

46. Please document if member has other behavioral health comorbidities (e.g., anxiety, depression, ADHD				

47. Please document medication trials with melatonin and/or clonidine, if clinically appropriate. Include drug name, dates/duration of use, and outcome.

Medication Name	Dates/Duration of Use	Outcomes
1.	1.	1.
2.	2.	2.
3.	3.	3.
4.	4.	4.

48.	Please document the clinical rationale for the use of a hypnotic agent for this member who is less than 6 years of age.		