


Vaccination ¹	Recommended Schedule
COVID-19 (<i>new 2022</i>)	Pfizer-BioNTech: 2 doses (21 days apart) Moderna: 2 doses (28 days apart) Janssen: 1 dose <i>(see next page for additional scheduling)</i>
Influenza inactivated (IIV4) or Influenza recombinant (RIV4)	1 dose annually (unless allergic to eggs)
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap (should have prior to turning 65), then Td or Tdap booster every 10 years
Zoster recombinant (RZV) (<i>preferred</i>) or Zoster live (ZVL) ²	2 doses (2-6 months apart) or 1 dose
Pneumococcal conjugate (PCV15)	1 dose (with PPSV23)
Pneumococcal polysaccharide (PPSV23)	1 dose (with PCV15)
Pneumococcal conjugate (PCV20) (<i>new 2022</i>) ³	1 dose
<ul style="list-style-type: none"> • If member has not had either Pneumococcal vaccine: <ul style="list-style-type: none"> ○ 1 dose PCV15, then 1 dose PPSV23 at least 1 year after ○ or 1 dose PCV20 • If member has had PPSV23, but not PCV15: <ul style="list-style-type: none"> ○ 1 dose PCV15 or PCV20 at least 1 year after last PPSV23 • If member has had PCV15, but not PPSV23: <ul style="list-style-type: none"> ○ 1 dose PPSV23 at least 1 year after last PCV15 	

Screening ⁴	Recommended Schedule
Mammogram ⁵	Every 2 years
Colonoscopy ⁶	Every 10 years (<i>preferred</i>) (<i>until age 75, then discuss with PCP whether to continue based on risk factors</i>)
 Other methods include: <ul style="list-style-type: none"> • High-sensitivity gFOBT or FIT every year • sDNA-FIT every 1 to 3 years 	<ul style="list-style-type: none"> • CT colonography every 5 years • Flexible sigmoidoscopy every 5 years • Flexible sigmoidoscopy every 10 years + FIT every year
Hearing exam	Every 2 years
Eye exam (members with Diabetes)	Dilated eye exam & Glaucoma screening annually

¹ CDC Vaccination updates – <https://www.cdc.gov/vaccines/schedules/hcp/imz/adult.html>

² CDC information: Shingles vaccination (zoster recombinant) – <https://www.cdc.gov/shingles/vaccination.html>

³ CDC information: Pneumococcal vaccinations – <https://www.cdc.gov/pneumococcal/vaccination.html>

⁴ US Preventive Services Task Force – <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation-topics/uspstf-a-and-b-recommendations>

⁵ CDC information: Breast cancer – <https://www.cdc.gov/cancer/breast/index.htm>

⁶ CDC information: Colorectal cancer screening tests – https://www.cdc.gov/cancer/colorectal/basic_info/screening/tests.htm

Screening ⁴	Recommended Schedule
Eye exam (members w/o Diabetes)	Dilated eye exam & Glaucoma screening every 2 years

COVID-19 Vaccination detail information⁷ – Specific to Age 65+

Information retrieved May 12, 2022 from Reference Guide published April 27, 2022. For potential updates: <https://www.cdc.gov/vaccines/covid-19/downloads/covid19-vaccine-quick-reference-guide-2pages.pdf>

Dose Type	Pfizer-BioNTech	Moderna	Janssen
Type of Vaccine	mRNA	mRNA	Viral vector
Primary Series	2 doses, separated by 21 days	2 doses, separated by 28 days	1 dose
Additional Primary Dose for severely immunocompromised persons	<ul style="list-style-type: none"> At least 28 days after second dose of Pfizer vaccine At least 28 days after one dose of Janssen vaccine 	<ul style="list-style-type: none"> At least 28 days after second dose of Moderna vaccine At least 28 days after one dose of Janssen vaccine 	Not authorized; Should receive dose of either mRNA vaccine
Booster schedule (general population)	<ul style="list-style-type: none"> At least 5 months after second primary dose of either mRNA vaccine At least 2 months after dose of Janssen vaccine 	<ul style="list-style-type: none"> At least 5 months after second primary dose of either mRNA vaccine At least 2 months after dose of Janssen vaccine 	** mRNA vaccines are preferred <ul style="list-style-type: none"> At least 2 months after primary dose of Janssen vaccine At least 3 months after second primary dose of either mRNA vaccine
Booster schedule (severely immunocompromised persons)	<ul style="list-style-type: none"> At least 3 months after additional primary dose of either mRNA vaccine At least 2 months after Janssen vaccine dose <u>and</u> additional dose of either mRNA vaccine 	<ul style="list-style-type: none"> At least 3 months after additional primary dose of either mRNA vaccine At least 2 months after Janssen vaccine dose <u>and</u> additional dose of either mRNA vaccine 	** mRNA vaccines are preferred <ul style="list-style-type: none"> At least 2 months after primary dose of Janssen vaccine At least 3 months after second primary dose of either mRNA vaccine

⁷ CDC information: US COVID-19 Vaccine Product Information – <https://www.cdc.gov/vaccines/covid-19/info-by-product/index.html>